

Gene D. Amman

Purchased by the Forest Service,
U. S. Department of Agriculture, for Official Use

Attraction of Mountain Pine Beetles¹ to Small-Diameter Lodgepole Pines Baited with *Trans-Verbenol* and *Alpha-Pinene*^{2,3}

LYNN A. RASMUSSEN⁴

USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah 84401

ABSTRACT

Lodgepole pines, *Pinus contorta* Douglas, 8.9 in. diameter at breast height (d.b.h.) or less, were baited with the pheromone *trans-verbenol* and the terpene *alpha-pinene* to determine if populations of *Dendroctonus ponderosae* Hopkins could be attracted to these trees of small diameter. Even though the beetles in most cases did not

successfully attack baited trees, such trees were usually the first to be attacked. This phenomenon indicated that these chemicals might be used to attract beetles into areas with baited trees but not to baited trees exclusively. This indication was supported by the fact that the beetles successfully attacked larger unbaited trees.

Infestations of mountain pine beetle, *Dendroctonus ponderosae* Hopkins, in lodgepole pine, *Pinus contorta* Douglas, have occurred in Idaho, Montana, and Utah at rather frequent intervals. The mountain pine beetle kills most of the large lodgepole pine

trees in a forest before the beetle population subsides, and it probably is the most aggressive *Dendroctonus* bark beetle in the western United States.

Amman (1969) reported that brood survival, as measured by emergence holes, was correlated closely with bark depth but varied with stand density and plot elevation. The greatest proportion of thick-barked trees was in the large-diameter classes. He hypothesized that ploem thickness was the critical

¹ Coleoptera: Scolytidae.
² Received for publication Oct. 13, 1971.
³ Mr. Walter E. Cole endorses and communicates.
⁴ Biological Technician.

factor. Cole and Amman (1969) indicated that large infestations of mountain pine beetle depend on the presence of large-diameter trees within a stand of lodgepole pine, and they concluded that growth of the beetle population is food-limited, i.e., survival and population size are correlated highly with the quantity of phloem within the infested tree. Therefore, if materials such as the pheromone *trans-verbenol* and the terpene *alpha-pinene* could be used to attract beetles to small-diameter thin-phloem trees, population reduction might be attained.

The occurrence of pheromones in *Dendroctonus* species is well known (McMullen and Atkins 1962, Vité 1962). Pheromones are the means of communication that cause mass attack and probably function to maintain the species integrity of bark beetles inhabiting the same host tree. Pitman et al. (1968) reported that *trans-verbenol*, a terpene alcohol isolated from the hindgut of the female, is the principal attractant for *D. ponderosae*. According to Pitman (1971), the effectiveness of *trans-verbenol* depended upon the terpene *alpha-pinene* produced by the host tree.

Even though the ability of *trans-verbenol* to attract the mountain pine beetle to white pine is substantiated, the question still remains whether *trans-verbenol* can be used to manipulate and manage populations of this beetle in lodgepole pine forests.

METHODS.—Two lodgepole pine stands in southeastern Idaho were selected in the spring of 1970. Each stand had a fairly uniform representation of 8- to 16-in.-diam trees and contained scattered groups of mountain pine beetle infestations. Nine groups of infested trees were designated as test plots; 8 were circular and 1 was rectangular.

Two plot designs were used. The 1st plot design consisted of 8 circular plots. For each of the circular plots, pairs of uninfested trees were selected around a group of infested trees at a radial distance of 198 ft (3 chains); the uninfested trees were so selected that they were a minimum of 66 ft (1 chain) apart. Each pair consisted of 1 tree 7- to 8.9-in. dbh with thin phloem (0.10 in. or less) and a 2nd tree 12 in. or larger dbh with thick phloem (0.11 in. or greater). The smaller was baited with *trans-verbenol* and *alpha-pinene*, whereas the larger tree was used as a check. The number of these pairs on the 8 circular plots ranged from 3 to 8, depending upon the avail-

ability of trees having desired characteristics. Each of the circular plots contained ca. 2.8 acres; thus, the total acreage involved was ca. 22.4 acres.

The 2nd plot design consisted of 1 rectangular plot. In this plot, 4 cruise lines were established 132 ft (2 chains) apart, and 45 baited trees alternating with 45 check trees were marked along these lines at 66-ft intervals. The size and phloem thickness requirements for baited and check trees were the same as those for the circular plots. This plot covered ca. 19 acres and contained 82 trees that had been infested by the beetle.

Using a method similar to that of Pitman (1971), 4-ft lengths of polyethylene catheter tubing (0.038 in. ID), charged with a mixture of *trans-verbenol* and *alpha-pinene* at a ratio of 9:1, were attached to each experimental tree. A polyethylene Boston bottle containing 10 ml of *alpha-pinene* also was attached to each of these trees to compensate for the higher evaporative rate of *alpha-pinene* from the mixture.

Daily observations were made to detect timing, sequence, and severity of beetle attacks. Also, attacks were recorded on nonexperimental trees within 66 ft of each baited tree in both types of plots.

RESULTS AND DISCUSSION.—The effectiveness of *trans-verbenol* and *alpha-pinene* in attracting mountain pine beetles is indicated by the relative proportions of trees attacked. Of the 64 trees that were attacked by the mountain pine beetle in the circular plots, 17% were baited, 5% were checks, and 78% were nonexperimental trees. Of the 84 trees that were attacked in the rectangular plot, 23% were baited, 2% were checks, and 75% were nonexperimental (Table 1). When only the baited and check trees were compared, the effectiveness of the attractants is indicated by the percentages of trees attacked: i.e., 22 and 6%, respectively, for the 8 circular plots; 42 and 4%, respectively, for the rectangular plot (Table 1).

Eleven of the 50 baited trees were attacked by the mountain pine beetle in the circular plots. However, only 6 (54%) of these were successfully mass attacked; the other 5 trees were considered to be attacked unsuccessfully because they either experienced only one to a few scattered attacks or the beetles were pitched out by the tree. Three of the 50 check trees were attacked; on 2 (66%) the attacks were successful. Fifty nonexperimental trees also were at-

Table 1.—Comparison of lodgepole pine trees attacked by the mountain pine beetle in circular and rectangular plots.

Plot	No. trees attacked			% trees attacked			% experimental trees attacked	
	Baited	Check	Non-experimental	Baited	Check	Non-experimental	Baited	Check
Circular								
1	1	1	11	8	8	84	14	14
2	1	0	4	20	0	80	17	0
3	1	0	6	14	0	86	33	0
4	2	0	12	14	0	86	29	0
5	3	1	9	23	8	69	43	14
6	0	0	2	0	0	100	0	0
7	2	1	5	25	12.5	62.5	33	17
8	1	0	1	50	0	50	14	0
Total	11	3	50	17	5	78	22	6
Rectangular	19	2	63	23	2	75	42	4




FIG. 1, 2.—Relationship of number of trees attacked by mountain pine beetle to time and distance from the nearest baited tree that was attacked. Fig. 1.—In the circular plots. Fig. 2.—In the rectangular plot.

tacked within these plots; on 46 trees (92%) the attacks were successful.

In the rectangular plot, 19 of the 45 baited trees were attacked but only 11 (58%) of these were attacked successfully. Two of the 45 check trees were attacked; on only 1 (50%) was the attack successful. Fifty-nine (94%) of 63 nonexperimental trees were attacked successfully.

In each of 5 circular plots, 1 baited tree was attacked first by the mountain pine beetle. In 1 circular plot, 1 baited, 1 check, and 1 nonexperimental tree were attacked at the same time; there was no way to distinguish which was attacked first. In 2 of the circular plots, nonexperimental trees were attacked first.

In the rectangular plot, 8 baited trees were attacked before any other tree. After 28 days of observation, 15 baited trees were attacked in contrast to

1 check tree and 9 nonexperimental trees. Thereafter, the number of nonexperimental trees being attacked rapidly increased until the beetle flight period was over.

In both the circular (Fig. 1) and rectangular (Fig. 2) plots, most of the attacked trees were in close proximity to the baited trees during the early part of the attack period. However, as the attack period progressed and the number of attacked trees increased, the proximal distance also increased.

A greater number of smaller trees, most of which were baited, were attacked and killed during the 1970 attractant test than had been killed in the previous year. In the 1970 attractant test, 20% of all the trees attacked within the circular plots and 32% within the rectangular plot were 8.9 in. or smaller (Table 2).

In the circular plots, the number of lodgepole

Table 2.—Comparison of lodgepole pine trees killed by the mountain pine beetle per year.

Plot	No. trees attacked	No. attacked 8.9 in. or less	% attacked 8.9 in. or less	Avg. DBH of trees attacked
Circular 1969	38	3	8	13.3
Circular 1970	64	13	20	13.0
Rectangular 1969	82	1	1	13.7
Rectangular 1970	84	27	32	11.8

pine trees attacked in 1970 was nearly double the number attacked in 1969, whereas in the rectangular plot the number remained almost equal for the 2 years (Table 2). A possible reason is that the general area of the circular plots contained a number of groups of trees infested by mountain pine beetles. Large numbers of the emerging beetles probably were attracted into the circular plots by the bait mixture and the newly attacked trees. However, in the rectangular plot, all the trees infested in 1969 were situated within the plot boundaries, and no supplemental source of beetles outside the plot was observed. Therefore, the number of trees attacked in 1970 more closely approximated the number attacked in 1969.

This study indicates the use of *trans-verbenol* and *alpha-pinene* to regulate mountain pine beetle populations by attracting beetles to small lodgepole pine where brood production is low does not appear possible at this time.

ACKNOWLEDGMENT.—The author thanks Dr. Gary B. Pitman of Boyce Thompson Institute for Plant Research for providing the *trans-verbenol*, *alpha-pinene* delivery system, and technical advice.

REFERENCES CITED

- Amman, G. D. 1969. Mountain pine beetle emergence in relation to depth of lodgepole pine bark. U.S. Forest Serv. Res. Note INT-96. 8 p.
- Colc, W. E., and G. D. Amman. 1969. Mountain pine beetle infestations in relation to lodgepole pine diameters. *Ibid.* 95: 7 p.
- McMullen, L. H., and M. D. Atkins. 1962. On the flight and host selection of the Douglas-fir beetle, *Dendroctonus pseudotsugae* Hopk. (Coleoptera: Scolytidae). *Can. Entomol.* 94: 1309-25.
- Pitman, G. B. 1971. *Trans-verbenol* and *alpha-pinene*: their utility in manipulation of the mountain pine beetle. *J. Econ. Entomol.* 64: 426-30.
- Pitman, G. B., J. P. Vité, G. W. Kinzer, and A. F. Fentiman, Jr. 1968. Bark beetle attractants: *trans-verbenol* isolated from *Dendroctonus*. *Nature (London)* 218: 168-9.
- Vité, J. P. 1962. The host factor in the population dynamics of bark beetles attacking ponderosa pine. *Proc. IUFRO, 13th Congr., Vienna, 1961, Part 2, vol. 1 (24-10)*. 8 p.