


CCAPS National Comparison — Pre-Post Change


VER. 5/2017

Based on 760 unique local clients, with at least 2 CCAPS Administrations from 07/01/2016 to 06/30/2017.
The clients in the resulting sample have an average of 6 administrations.

Change for Clients with Moderate and/or Elevated Distress



Change for Clients with Elevated Distress



This report compares your center's average change on the CCAPS' subscales to a national sample of 106 counseling centers representing 47,948 clients. Average change is calculated by subtracting each client's last scorible CCAPS administration from their first scorible CCAPS administration and then averaging these differences by subscale. Two charts are provided above based on client's level of initial distress: (1) clients with moderate and/or elevated initial distress or (2) clients with only elevated initial distress.

For example, looking at the first chart (moderate + elevated), your center's average change for Depression is at the 52nd percentile. This means that your center's average change on the Depression subscale is greater than the change achieved by 52 % of counseling centers in the national sample (for clients who's initial distress was at least moderate and/or elevated).

CCAPS Pre-Post Change Data Table

Subscale by Initial Distress Level	Local Average Change in Raw Subscale Score	# of Local Clients Above Cut Score	% of Local Clients Above Cut Score	National Average Change in Raw Subscale Score	# of National Clients Above Cut Score	% of National Clients above Cut Score
Depression Low-Cut	0.686	627	82.5 %	0.673	34,356	71.7 %
Depression High-Cut	0.852	454	59.7 %	0.835	24,369	50.8 %
Generalized Anxiety Low-Cut	0.555	586	77.1 %	0.523	35,923	74.9 %
Generalized Anxiety High-Cut	0.673	382	50.3 %	0.684	22,916	47.8 %
Social Anxiety Low-Cut	0.452	544	71.6 %	0.419	30,170	62.9 %
Social Anxiety High-Cut	0.553	339	44.6 %	0.525	17,372	36.2 %
Academic Distress Low-Cut	0.405	549	72.2 %	0.475	32,448	67.7 %
Academic Distress High-Cut	0.655	295	38.8 %	0.776	14,468	30.2 %
Eating Concerns Low-Cut	0.588	276	36.3 %	0.595	16,463	34.3 %
Eating Concerns High-Cut	0.653	221	29.1 %	0.662	13,343	27.8 %
Hostility Low-Cut	0.625	363	47.8 %	0.545	22,581	47.1 %
Hostility High-Cut	0.847	214	28.2 %	0.780	11,604	24.2 %
Alcohol Use Low-Cut	0.644	130	17.1 %	0.570	15,870	33.1 %
Alcohol Use High-Cut	0.843	86	11.3 %	0.749	10,163	21.2 %
Distress Index Low-Cut	0.548	588	77.4 %	0.490	36,138	75.4 %
Distress Index High-Cut	0.656	332	43.7 %	0.672	16,856	35.2 %

Data Table Column Descriptions

Subscale by Initial Distress Level	Scores on the CCAPS are divided into three ranges of distress, Low-Distress (white), Moderate Distress (yellow), and Elevated Distress (red), using Low and High cut-scores (Low-Cut and High-Cut). Clients that score over the Low-Cut include Moderate Distress plus Elevated Distress scores. Those that score over the High-Cut include Elevated Distress scores only. For more information about cut-scores please view the CCAPS User Manual under "CCAPS" in the Help Tab.
Local Average Change in Raw Subscale Score	The average difference in Pre-Post CCAPS scores at your center for a given subscale and initial distress.
# of Local Clients Above Cut Score	The number of clients at your center that scored above the indicated cut-score (low or high) per subscale.
% of Local Clients Above Cut Score	The percentage of clients at your center that scored the indicated cut-score (low or high) per subscale.
National Average Change in Raw Subscale Score	The national average Pre-Post change in CCAPS scores for the indicated subscale and cut-score.
# of National Clients Above Cut Score	The number of clients in the national sample that scored above the indicated subscale and cut-score.
% of National Clients Above Cut Score	The percentage of clients in the national sample that scored above the indicated subscale and cut-score.