

## Math 4200

**Theorem.** Every convergent sequence of real numbers is bounded.

proof:

Let  $\{a_n\}$  be a sequence and suppose  $a_n \rightarrow L$ . For  $\epsilon = 13$ , there exists a positive integer  $M > 0$  such that if  $n > M$  then  $|a_n - L| < 13$ . That is, if  $n > M$  then  $L - 13 < a_n < L + 13$ .

Now let  $B = \max\{|a_1|, |a_2|, \dots, |a_M|, |L - 13|, |L + 13|\}$ . If  $1 \leq n \leq M$ , then clearly  $|a_n| \leq B$ . If  $n > M$  then  $L - 13 < a_n < L + 13$ , and so  $|a_n| \leq \max\{|L - 13|, |L + 13|\}$ . Hence, for all  $n$ ,  $|a_n| \leq B$ . Therefore,  $\{a_n\}$  is bounded.

**Theorem.** If  $\{a_n\}$  and  $\{b_n\}$  are sequences having limits  $A$  and  $B$  respectively, then the sequence  $\{a_n \cdot b_n\}$  converges to  $A \cdot B$ .

proof:

Since  $\{a_n\}$  is bounded, there exists  $K > 0$  such that  $|a_n| \leq K$  for each  $n$ . Let  $\epsilon > 0$ . Since  $\{b_n\}$  converges to  $B$ , there exists  $M_1 > 0$  such that if  $n > M_1$  then  $|b_n - B| < \frac{\epsilon}{2K}$ . Since  $\{a_n\}$  converges to  $A$ , there exists  $M_2 > 0$  such that if  $n > M_2$  then  $|a_n - A| < \frac{\epsilon}{2(|B|+1)}$ . Choose  $M = \max\{M_1, M_2\}$ . For  $n > M$ ,  
 $|a_n b_n - AB| = |a_n b_n - a_n B + a_n B - AB| \leq |a_n b_n - a_n B| + |a_n B - AB| = |a_n| |b_n - B| + |B| |a_n - A| < K \cdot \frac{\epsilon}{2K} + |B| \cdot \frac{\epsilon}{2(|B|+1)} < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$ . Hence, the sequence  $\{a_n \cdot b_n\}$  converges to  $A \cdot B$ .